arXiv:1208.0879v2 [quant-ph] 27 Aug 2012

Quantum Deficit and Geometry for a Class of Two-qubit States

Yao-Kun Wang,[1, 2] Teng Ma,[1] Bo Li,[3, 4] and Zhi-Xi Wang[1]
[1]*School of Mathematical Sciences, Capital Normal University, Beijing 100048, China*
[2]*Department of Mathematics, Tonghua Normal University, Tonghua 134001, China*
[3]*Institute of Physics, Chinese Academy of Sciences, Beijing 100190, China*
[4]*Department of Mathematics and Computer,*
Shangrao Normal University, Shangrao 334001, China

Quantum deficit, as introduced by Jonathan Oppenheim *et al* [Phys. Rev. Lett. **89**, 180402 (2002)] is a good measure of the quantum correlations in a state and provides a new form for understanding quantum nonlocality, and is different from quantum discord. We will analytically evaluate the quantum deficit of the Bell-diagonal states and a class of two-qubit states and give the geometry of those quantum deficit. The dynamic behavior of quantum deficit under decoherence is investigated and it is shown that the quantum deficit of some classes of X states is more robust against the decoherence than the entanglement.

I. INTRODUCTION

The quantum correlation of quantum states includes entanglement and other kinds of nonclassical correlation, and the quantum correlations is more general than entanglement [1, 2]. A formal measure of quantum correlations in measurements is quantum discord [2]. Quantum discord describes a different aspect of nonclassical correlation. Many works have been done to the importance and application of quantum discord. In particular, there is accurate expression for quantum discord for two-qubit states, even for the X states [3–5], and the geometry of the quantum correlation about those states is introduced by [6–9]. Recently, researches on the dynamics of quantum discord in various noisy environments have revealed many fascinating features [10], and manifested that discord is more robust than entanglement for both Markovian and non-Markovian dissipative processes, and moreover, after a phase damping process, the quantum discord of a type of Bell-diagonal state was shown to exhibit a freezing phenomenon.

There are also a lot of measures of nonclassical correlation besides entanglement, such as the quantum deficit [11, 12], measurement-induced disturbance [13], symmetric discord [14, 15], relative entropy of discord and dissonance [16],geometric discord [7, 8], and continuous-variable discord [17, 18], and so on. Almost all different measures of quantum correlation beyond entanglement were reviewed in [19]. The work deficit[11] is the first operational approach to quantify quantum correlations. Then, a similar process was used to justify a physical interpretation of quantum discord [20]. In the physical view, the results of [11] showed that quantum deficit originates in questions using nonlocal operation to extract work from a correlated system coupled to a heat bath only in the case of pure states, and in the general case, the advantage is related to more general forms of quantum correlations. The corresponding notion of classicality is in the spirit of LOCC [12]. Jonathan Oppenheim *et al* [11] define quantum deficit as the work deficit

$$\Delta \equiv W_t - W_l. \tag{1}$$

Horodecki *et al* interpret that quantum deficit is the difference between the information of the whole system and the localizable information and give various simpler forms of deficit [21]. Synak *et al* prove that the deficit is lower-bounded by the (regularized) relative entropy of entanglement for werner and isotropic states

[22]. Recently, Alexander Streltsov *et al* give the definition of one-way information deficit by the relative entropy over all local vonNeumann measurements on oneside and reveal the fundamental role of quantum correlations as a resource for the distribution of entanglement [23, 24], and also give the other definition of one-way information deficit by vonNeumann measurements on oneside [25]

$$\Delta^{\rightarrow}(\rho^{ab}) = \min_{\{\Pi_k\}} S(\sum_i \Pi_k \rho^{ab} \Pi_k) - S(\rho^{ab}). \tag{2}$$

Quantum discord and quantum deficit is different types of quantum correlation. We have known that the geometry and the dynamics of quantum discord for a class of two-qubit states by account above , and analogously, how about their quantum deficit? In this article, we will give the answer.

Undergoing proper local unitary transformations, any two-qubit state ρ^{ab} can be written as:

$$\rho^{ab} = \frac{1}{4}(I \otimes I + \mathbf{r} \cdot \sigma \otimes I + I \otimes \mathbf{s} \cdot \sigma + \sum_{i=1}^{3} c_i \sigma_i \otimes \sigma_i), \tag{3}$$

where $\mathbf{r}$ and $\mathbf{s}$ are Bloch vectors and $\{\sigma_i\}_{i=1}^3$ are the standard Pauli matrices. When $\mathbf{r}$=$\mathbf{s}$=$\mathbf{0}$, ρ reduces to the two-qubit Bell-diagonal states. Next, we assume that the Bloch vectors are z directional, that is, $\mathbf{r} = (0, 0, r)$, $\mathbf{s} = (0, 0, s)$. We can also change them to be x or y directional via an appropriate local unitary transformation without losing its diagonal property of the correlation terms [26]. In this case a state ρ defined in Eq.(3) has the form

$$\rho^{ab} = \frac{1}{4}\begin{pmatrix} 1+r+s+c_3 & 0 & 0 & c_1-c_2 \\ 0 & 1+r-s-c_3 & c_1+c_2 & 0 \\ 0 & c_1+c_2 & 1-r+s-c_3 & 0 \\ c_1-c_2 & 0 & 0 & 1-r-s+c_3 \end{pmatrix}. \tag{4}$$

We calculate the quantum deficit defined by Eq.(2) of Bell-Diagonal states and point out that the quantum deficit of Bell-Diagonal states equal to its quantum discord in sec. II. We depict the level surface of constant deficit in four different situations in sec. III. In sec. IV, we discuss the dynamics of quantum deficit and show that the quantum deficit of a certain class of X states decay under decoherence. A brief conclusion is given in sec. V.

II. THE DEFICIT OF BELL-DIAGONAL STATES

For the two-qubit Bell-diagonal state:

$$\rho^{ab} = \frac{1}{4}(I \otimes I + \sum_{i=1}^{3} c_i \sigma_i \otimes \sigma_i). \tag{5}$$

The eigenvalues of ρ are

$$\lambda_1 = \frac{1}{4}(1 - c_1 - c_2 - c_3), \lambda_2 = \frac{1}{4}(1 - c_1 + c_2 + c_3), \lambda_3 = \frac{1}{4}(1 + c_1 - c_2 + c_3), \lambda_4 = \frac{1}{4}(1 + c_1 + c_2 - c_3).$$

The entropy of ρ^{ab} is

$$\begin{aligned} S(\rho^{ab}) &= -\sum_{i=1}^{4} \lambda_i \log \lambda_i \\ &= 2 - \frac{1-c_1-c_2-c_3}{4}\log(1-c_1-c_2-c_3) \\ &\quad - \frac{1-c_1+c_2+c_3}{4}\log(1-c_1+c_2+c_3) \\ &\quad - \frac{1+c_1-c_2+c_3}{4}\log(1+c_1-c_2+c_3) \\ &\quad - \frac{1+c_1+c_2-c_3}{4}\log(1+c_1+c_2-c_3). \end{aligned} \tag{6}$$

Next, we evaluate the the quantum deficit. Let

$$\{\Pi_k = |k\rangle\langle k|, k = 0, 1\} \tag{7}$$

be the local measurement for the part b along the computational base $|k\rangle$; then any von Neumann measurement for the part b can be written as

$$\{B_k = V\Pi_k V^\dagger : k = 0, 1\} \tag{8}$$

for some unitary $V \in U(2)$. But any unitary V can be written, up to a constant phase, as

$$V = tI + i\overrightarrow{y}\overrightarrow{\sigma} \tag{9}$$

with $t \in R$, $\overrightarrow{y} = (y_1, y_2, y_3) \in R^3$, and $t^2 + y_1^2 + y_2^2 + y_3^2 = 1$. After the measurement B_k, the state ρ^{ab} will change to the ensemble ρ_k, p_k with

$$\rho_k := \frac{1}{p_k}(I \otimes B_k)\rho(I \otimes B_k) \tag{10}$$

and $p_k = tr(I \otimes B_k)\rho(I \otimes B_k)$. We need to evaluate ρ_k and p_k. For this purpose, we write

$$p_k\rho_k = (I \otimes B_k)\rho(I \otimes B_k) = \frac{1}{4}(I \otimes V)(I \otimes \Pi_k)(I + \sum_{j=1}^{3} c_j\sigma_j \otimes (V^\dagger\sigma_j V))(I \otimes \Pi_k)(I \otimes V^\dagger). \tag{11}$$

By use of the relations [13]

$$V^\dagger\sigma_1 V = (t^2 + y_1^2 - y_2^2 - y_3^2)\sigma_1 + 2(ty_3 + y_1y_2)\sigma_2 + 2(-ty_2 + y_1y_3)\sigma_3, \tag{12}$$

$$V^\dagger\sigma_2 V = 2(-ty_3 + y_1y_2)\sigma_1 + (t^2 + y_2^2 - y_1^2 - y_3^2)\sigma_2 + 2(ty_1 + y_2y_3)\sigma_3, \tag{13}$$

$$V^\dagger\sigma_3 V = 2(ty_2 + y_1y_3)\sigma_1 + 2(-ty_1 + y_2y_3)\sigma_2 + (t^2 + y_3^2 - y_1^2 - y_2^2)\sigma_3, \tag{14}$$

and

$$\Pi_0\sigma_3\Pi_0 = \Pi_0, \Pi_1\sigma_3\Pi_1 = -\Pi_1, \Pi_j\sigma_k\Pi_j = 0, forj = 0, 1, k = 1, 2. \tag{15}$$

We obtain $p_0 = p_1 = \frac{1}{2}$ and

$$\rho_0 = \frac{1}{2}(I + c_1z_1\sigma_1 + c_2z_2\sigma_2 + c_3z_3\sigma_3) \otimes (V\Pi_0V^\dagger), \tag{16}$$

$$\rho_1 = \frac{1}{2}(I - c_1z_1\sigma_1 - c_2z_2\sigma_2 - c_3z_3\sigma_3) \otimes (V\Pi_1V^\dagger), \tag{17}$$

where

$$z_1 = 2(-ty_2 + y_1y_3), z_2 = 2(ty_1 + y_2y_3), z_3 = t^2 + y_3^2 - y_1^2 - y_2^2. \tag{18}$$

Let $M = c_1z_1\sigma_1 + c_2z_2\sigma_2 + c_3z_3\sigma_3$, then

$$\rho_0 = \frac{1}{2}(I + M) \otimes (V\Pi_0V^\dagger), \rho_1 = \frac{1}{2}(I - M) \otimes (V\Pi_1V^\dagger), \tag{19}$$

The eigenvalues of $\frac{1}{2}(I+M)$ and $\frac{1}{2}(I-M)$ are $\frac{1-\theta}{2}$, $\frac{1+\theta}{2}$ and $\frac{1+\theta}{2}$, $\frac{1-\theta}{2}$, where $\theta = \sqrt{|c_1z_1|^2 + |c_2z_2|^2 + |c_3z_3|^2}$. We can check that $\frac{1}{2}(I + M)$ commutes with $\frac{1}{2}(I - M)$, and there exists an orthogonal basis such that both $\frac{1}{2}(I + M)$ and $\frac{1}{2}(I - M)$ are diagonal with respect to that basis [27], i.e for some unitary $U \in U(2)$, and

$$\frac{1}{2}(I + M) = U \begin{pmatrix} \frac{1-\theta}{2} & 0 \\ 0 & \frac{1+\theta}{2} \end{pmatrix} U^\dagger, \frac{1}{2}(I - M) = U \begin{pmatrix} \frac{1+\theta}{2} & 0 \\ 0 & \frac{1-\theta}{2} \end{pmatrix} U^\dagger. \tag{20}$$

As $\sum_i \Pi_k\rho^{ab}\Pi_k = p_0\rho_0 + p_1\rho_1$, and then we evaluate the eigenvalues of $\sum_i \Pi_k\rho^{ab}\Pi_k$,

$$\begin{aligned}
& |\lambda E - \sum_i \Pi_k\rho^{ab}\Pi_k| \\
= \; & |\lambda E - (p_0\rho_0 + p_1\rho_1)| \\
= \; & \left|\lambda E - (\frac{1}{2}\rho_0 + \frac{1}{2}\rho_1)\right| \\
= \; & \left|\lambda E - \{[U \begin{pmatrix} \frac{1-\theta}{4} & 0 \\ 0 & \frac{1+\theta}{4} \end{pmatrix} U^\dagger] \otimes (V\Pi_0V^\dagger) + [U \begin{pmatrix} \frac{1+\theta}{4} & 0 \\ 0 & \frac{1-\theta}{4} \end{pmatrix} U^\dagger] \otimes (V\Pi_1V^\dagger)\}\right| \\
= \; & \left|\lambda E - \{(U \otimes V)[\begin{pmatrix} \frac{1-\theta}{4} & 0 \\ 0 & \frac{1+\theta}{4} \end{pmatrix}] \otimes \Pi_0](U^\dagger \otimes V^\dagger) + \{(U \otimes V)[\begin{pmatrix} \frac{1+\theta}{4} & 0 \\ 0 & \frac{1-\theta}{4} \end{pmatrix}] \otimes \Pi_1](U^\dagger \otimes V^\dagger)\}\right| \\
= \; & \left|\lambda E - (U \otimes V)[\begin{pmatrix} \frac{1-\theta}{4} & 0 \\ 0 & \frac{1+\theta}{4} \end{pmatrix} \otimes \Pi_0 + \begin{pmatrix} \frac{1+\theta}{4} & 0 \\ 0 & \frac{1-\theta}{4} \end{pmatrix} \otimes \Pi_1](U^\dagger \otimes V^\dagger)\right| \\
= \; & \left|(U \otimes V)\lambda E(U \otimes V)^\dagger - (U \otimes V) \begin{pmatrix} \frac{1-\theta}{4} & 0 & 0 & 0 \\ 0 & \frac{1+\theta}{4} & 0 & 0 \\ 0 & 0 & \frac{1+\theta}{4} & 0 \\ 0 & 0 & 0 & \frac{1-\theta}{4} \end{pmatrix} (U \otimes V)^\dagger\right| \\
= \; & |(U \otimes V)| \cdot \left|\lambda E - \begin{pmatrix} \frac{1-\theta}{4} & 0 & 0 & 0 \\ 0 & \frac{1+\theta}{4} & 0 & 0 \\ 0 & 0 & \frac{1+\theta}{4} & 0 \\ 0 & 0 & 0 & \frac{1-\theta}{4} \end{pmatrix}\right| \cdot |(U \otimes V)^\dagger| \\
= \; & \left|\lambda E - \begin{pmatrix} \frac{1-\theta}{4} & 0 & 0 & 0 \\ 0 & \frac{1+\theta}{4} & 0 & 0 \\ 0 & 0 & \frac{1+\theta}{4} & 0 \\ 0 & 0 & 0 & \frac{1-\theta}{4} \end{pmatrix}\right|.
\end{aligned} \tag{21}$$

And the eigenvalues of $\sum_i \Pi_k\rho^{ab}\Pi_k$ are $\lambda_5 = \frac{1-\theta}{4}$, $\lambda_6 = \frac{1+\theta}{4}$, $\lambda_7 = \frac{1+\theta}{4}$, $\lambda_8 = \frac{1-\theta}{4}$. The entropy of $\sum\limits_i \Pi_k\rho^{ab}\Pi_k$ is

$$\begin{aligned}
S(\sum_i \Pi_k\rho^{ab}\Pi_k) &= -\sum_{i=5}^{8} \lambda_i \log \lambda_i \\
&= 2 - \frac{1-\theta}{2}\log(1-\theta) - \frac{1+\theta}{2}\log(1+\theta).
\end{aligned} \tag{22}$$

It can be directly verified that

$$z_1^2 + z_2^2 + z_3^2 = 1. \tag{23}$$

Let us put $c := \max\{|c_1|, |c_2|, |c_3|\}$, then $\theta \leq \sqrt{|c|^2(|z_1|^2+|z_2|^2+|z_3|^2)} = c$. Therefore, we see that

$$\sup_{\{B_k\}} \theta = \sup_{\{V\}} \theta = c. \tag{24}$$

The range of values allowed for θ is $\theta \in [0, c]$, and the derivative of $S(\sum_i \Pi_k \rho^{ab} \Pi_k)$ is $\frac{1}{2}[\log(1-\theta) - \log(1+\theta)]$, and it can be verified that it is monotonic decreasing function in the interval of $[0, c]$, and the minimal value of $S(\sum_i \Pi_k \rho^{ab} \Pi_k)$ can be attained at point c,

$$\min_{\{\Pi_k\}} S(\sum_i \Pi_k \rho^{ab} \Pi_k) = 2 - \frac{1-c}{2}\log(1-c) - \frac{1+c}{2}\log(1+c). \tag{25}$$

By Eq.(6), (25), and the quantum deficit of ρ^{ab} is given by

$$\begin{aligned} \Delta^{\rightarrow}(\rho^{ab}) &= \min_{\{\Pi_k\}} S(\sum_i \Pi_k \rho^{ab} \Pi_k) - S(\rho^{ab}) \\ &= \frac{1}{4}[(1-c_1-c_2-c_3)\log(1-c_1-c_2-c_3) \\ &\quad +(1-c_1+c_2+c_3)\log(1-c_1+c_2+c_3) \\ &\quad +(1+c_1-c_2+c_3)\log(1+c_1-c_2+c_3) \\ &\quad +(1+c_1+c_2-c_3)\log(1+c_1+c_2-c_3)] \\ &\quad -\frac{1-c}{2}\log(1-c) - \frac{1+c}{2}\log(1+c). \end{aligned} \tag{26}$$

Notice that the above result about the quantum deficit of ρ^{ab} coincides with that of quantum discord obtained in [13]. The geometry picture of the quantum discord given in Figure 2 of [6].

III. GEOMETRICAL DEPICTION OF A CLASS OF X STATES

For the state in Eq.(4), let $r = 0$, we obtain the following state:

$$\rho^{ab} = \frac{1}{4}\begin{pmatrix} 1+s+c_3 & 0 & 0 & c_1-c_2 \\ 0 & 1-s-c_3 & c_1+c_2 & 0 \\ 0 & c_1+c_2 & 1+s-c_3 & 0 \\ c_1-c_2 & 0 & 0 & 1-s+c_3 \end{pmatrix}, \tag{27}$$

and we will only consider the following further simplified family of the Eq.(27), where

$$|c_1| < |c_2| < |c_3|, 0 < |s| < 1 - |c_3|. \tag{28}$$

The concurrence of the state in Eq.(27), (28) can be calculated in terms of the eigenvalues of $\rho\widetilde{\rho}$, where $\widetilde{\rho} = \sigma_y \otimes \sigma_y \rho^* \sigma_y \otimes \sigma_y$. For the state, the eigenvalues of $\rho\widetilde{\rho}$ are

$$\begin{aligned} \lambda_9 &= \frac{1}{16}(c_1 - c_2 - \sqrt{(1+c_3)^2 - s^2})^2 \\ &= \frac{1}{16}(c_1 - c_2 - \sqrt{(1+s+c_3)(1-s+c_3)})^2, \end{aligned}$$

$$\begin{aligned} \lambda_{10} &= \frac{1}{16}(c_1 - c_2 + \sqrt{(1+c_3)^2 - s^2})^2 \\ &= \frac{1}{16}(c_1 - c_2 + \sqrt{(1+s+c_3)(1-s+c_3)})^2, \end{aligned}$$

$$\begin{aligned}\lambda_{11} &= \frac{1}{16}(c_1+c_2-\sqrt{(1-c_3)^2-s^2})^2\\ &= \frac{1}{16}(c_1+c_2-\sqrt{(1-s-c_3)(1+s-c_3)})^2,\end{aligned}$$

$$\begin{aligned}\lambda_{12} &= \frac{1}{16}(c_1+c_2+\sqrt{(1-c_3)^2-s^2})^2\\ &= \frac{1}{16}(c_1+c_2+\sqrt{(1-s-c_3)(1+s-c_3)})^2.\end{aligned}$$

The concurrence of the state in Eq.(27), (28) is given by

$$C(\rho^{ab})=\max\{2\max\{\sqrt{\lambda_9},\sqrt{\lambda_{10}},\sqrt{\lambda_{11}},\sqrt{\lambda_{12}}\}-\sqrt{\lambda_9}-\sqrt{\lambda_{10}}-\sqrt{\lambda_{11}}-\sqrt{\lambda_{12}},0\}. \tag{29}$$

Next, we will only consider the quantum deficit of the state in Eq.(27), (28). The computational methods of the others is similar to this situation, it doesn't merit any additional discussion here. Our computational procedure of the deficit is similar to the Bell-Diagonal case. In this case the eigenvalues of the state in Eq.(27), (28) are give by

$$\lambda_{13,14}=\frac{1}{4}[1-c_3\pm\sqrt{s^2+(c_1+c_2)^2}],\quad \lambda_{15,16}=\frac{1}{4}[1+c_3\pm\sqrt{s^2+(c_1-c_2)^2}].$$

The entropy is given by

$$\begin{aligned}S(\rho^{ab}) &= -\sum_{i=13}^{16}\lambda_i\log\lambda_i\\ &= 2-\frac{1}{4}[(1-c_3+\sqrt{s^2+(c_1+c_2)^2})\log(1-c_3+\sqrt{s^2+(c_1+c_2)^2})\\ &\quad +(1-c_3-\sqrt{s^2+(c_1+c_2)^2})\log(1-c_3-\sqrt{s^2+(c_1+c_2)^2})\\ &\quad +(1+c_3+\sqrt{s^2+(c_1+c_2)^2})\log(1+c_3+\sqrt{s^2+(c_1+c_2)^2})\\ &\quad +(1+c_3-\sqrt{s^2+(c_1+c_2)^2})\log(1+c_3-\sqrt{s^2+(c_1+c_2)^2})].\end{aligned} \tag{30}$$

Next, we will evaluate

$$\begin{aligned}p_k\rho_k &= (I\otimes B_k)\rho(I\otimes B_k)\\ &= \frac{1}{4}(I\otimes V)(I\otimes\Pi_k)(I\otimes V^\dagger)(I\otimes I+I\otimes s\sigma_3+\sum_{i=1}^{3}c_i\sigma_i\otimes\sigma_i))(I\otimes V)(I\otimes\Pi_k)(I\otimes V^\dagger)\\ &= \frac{1}{4}I\otimes V\Pi_kV^\dagger+\frac{s}{4}I\otimes V\Pi_kV^\dagger\sigma_3V\Pi_kV^\dagger+\sum_{i=1}^{3}\frac{c_i}{4}\sigma_i\otimes V\Pi_kV^\dagger\sigma_iV\Pi_kV^\dagger.\end{aligned} \tag{31}$$

By use of the Eq.(12), (13), (14), (15), (18), we obtain

$$p_0\rho_0=\frac{1}{4}(I+sz_3I+c_1z_1\sigma_1+c_2z_2\sigma_2+c_3z_3\sigma_3)\otimes V\Pi_0V^\dagger, \tag{32}$$

$$p_1\rho_1=\frac{1}{4}(I-sz_3I-c_1z_1\sigma_1-c_2z_2\sigma_2-c_3z_3\sigma_3)\otimes V\Pi_0V^\dagger. \tag{33}$$

Let $\overline{M}=sz_3I+c_1z_1\sigma_1+c_2z_2\sigma_2+c_3z_3\sigma_3$, and

$$p_0\rho_0=\frac{1}{4}(I+\overline{M})\otimes V\Pi_0V^\dagger,\quad p_1\rho_1=\frac{1}{4}(I-\overline{M})\otimes V\Pi_0V^\dagger. \tag{34}$$

Similar to Eq.(21), the eigenvalues of $\frac{1}{4}(I+\overline{M})$ and $\frac{1}{4}(I-\overline{M})$ are $\lambda_{17}=\frac{1}{4}(1+\phi-\theta), \lambda_{18}=\frac{1}{4}(1+\phi+\theta)$ and $\lambda_{19}=\frac{1}{4}(1-\phi-\theta), \lambda_{20}=\frac{1}{4}(1-\phi+\theta)$, where

$$\phi=sz_3, \theta=\sqrt{|c_1z_1|^2+|c_2z_2|^2+|c_3z_3|^2}. \tag{35}$$

The entropy of $\sum\limits_i \Pi_k\rho^{ab}\Pi_k$ is

$$\begin{aligned} S(\sum_i \Pi_k\rho^{ab}\Pi_k) &= f(\phi,\theta)=-\sum_{i=17}^{20}\lambda_i\log\lambda_i \\ &= 2-\frac{1}{4}[(1+\phi-\theta)\log(1+\phi-\theta)+(1+\phi+\theta)\log(1+\phi+\theta) \\ &\quad +(1-\phi-\theta)\log(1-\phi-\theta)+(1-\phi+\theta)\log(1-\phi+\theta)]. \end{aligned} \tag{36}$$

Using the domain of logarithmic function in $f(\phi,\theta)$ and Eq.(28) we obtain the range of θ and ϕ:

$$0\le|c_1|\le\theta\le|c_3|\le 1, -1<\phi<1. \tag{37}$$

We can verify that $f(-\phi,\theta)=f(\phi,\theta)$, the graph of $f(\phi,\theta)$ is symmetrical with respect to the θ-axis; $\frac{\partial f}{\partial\theta}=-\frac{1}{4}\log[\frac{(1+\theta)^2-\phi^2}{(1-\theta)^2-\phi^2}]<0, (0<\theta<1)$, $f(\phi,\theta)$ is a monotonic decreasing function in the positive direction of θ; $\frac{\partial f}{\partial\phi}=-\frac{1}{4}\log[\frac{(1+\phi)^2-\theta^2}{(1-\phi)^2-\theta^2}]<0, (0<\phi<1)$, $f(\phi,\theta)$ is a monotonic decreasing function in the positive direction of ϕ. When $\theta=|c_3|$, by Eq.(23), (28), (35) and we can obtain

$$\phi=|s|. \tag{38}$$

By Eq.(28), the projection of $f(\phi,\theta)$ on the plane $\phi o\theta$ is a symmetrical rectangle with respect to the θ-axis, and by use of the monotonicity of $f(\phi,\theta)$ in the positive direction of θ and ϕ, $f(\phi,\theta)$ can obtain the minimum in the point $(|s|,|c_3|)$, the minimum of $f(\phi,\theta)$ is given by

$$\begin{aligned} \min S(\sum_i \Pi_k\rho^{ab}\Pi_k) &= 2-\frac{1}{4}[(1+s-c_3)\log(1+s-c_3)+(1+s+c_3)\log(1+s+c_3) \\ &\quad +(1-s-c_3)\log(1-s-c_3)+(1-s+c_3)\log(1-s+c_3)]. \end{aligned} \tag{39}$$

By Eq.(30), (39), the quantum deficit of the state in Eq.(27), (28) is given by

$$\begin{aligned} \Delta^{\rightarrow}(\rho^{ab}) &= \min_{\{\Pi_k\}} S(\sum_i \Pi_k\rho^{ab}\Pi_k)-S(\rho^{ab}) \\ &= \frac{1}{4}[(1-c_3+\sqrt{s^2+(c_1+c_2)^2})\log(1-c_3+\sqrt{s^2+(c_1+c_2)^2}) \\ &\quad +(1-c_3-\sqrt{s^2+(c_1+c_2)^2})\log(1-c_3-\sqrt{s^2+(c_1+c_2)^2}) \\ &\quad +(1+c_3+\sqrt{s^2+(c_1+c_2)^2})\log(1+c_3+\sqrt{s^2+(c_1+c_2)^2}) \\ &\quad +(1+c_3-\sqrt{s^2+(c_1+c_2)^2})\log(1+c_3-\sqrt{s^2+(c_1+c_2)^2})] \\ &\quad -\frac{1}{4}[(1+s-c_3)\log(1+s-c_3)+(1+s+c_3)\log(1+s+c_3) \\ &\quad +(1-s-c_3)\log(1-s-c_3)+(1-s+c_3)\log(1-s+c_3)]. \end{aligned} \tag{40}$$

In Fig.1 we plot the level surface of deficit when (a) $s=0.3$, $\Delta^{\rightarrow}(\rho^{ab})=0.03$; (b) $s=0.5$, $\Delta^{\rightarrow}(\rho^{ab})=0.03$; (c) $s=0.3$, $\Delta^{\rightarrow}(\rho^{ab})=0.15$; (d) $s=0.5$, $\Delta^{\rightarrow}(\rho^{ab})=0.15$. From Fig.1 one can see that the level surface of deficit has a great change from the case $r = s = 0$ studied in Figure 2 of [6]. The surface shrinks with the effect s and the shrinking rate becomes larger with the increasing $|s|$. What is more, for larger deficit and small s (see Fig.(c)), the figure is similar to the ones in case of $r = s = 0$. But for larger s (see Fig.(d)), the figure is moved up again and changes dramatically also.



FIG. 1: Surfaces of constant deficit: (a) $s = 0.3$, $\Delta^{\rightarrow}(\rho^{ab}) = 0.03$; (b) $s = 0.5$, $\Delta^{\rightarrow}(\rho^{ab}) = 0.03$; (c) $s = 0.3$, $\Delta^{\rightarrow}(\rho^{ab}) = 0.15$; (d) $s = 0.5$, $\Delta^{\rightarrow}(\rho^{ab}) = 0.15$.

IV. DYNAMICS OF QUANTUM DEFICIT UNDER LOCAL NONDISSIPATIVE CHANNELS

In the following we consider that the state in Eq.(27), (28) undergoes the phase flip channel [28], with the Kraus operators $\Gamma_0^{(A)} = \mathrm{diag}(\sqrt{1-p/2}, \sqrt{1-p/2}) \otimes I$, $\Gamma_1^{(A)} = \mathrm{diag}(\sqrt{p/2}, -\sqrt{p/2}) \otimes I$, $\Gamma_0^{(B)} = I \otimes \mathrm{diag}(\sqrt{1-p/2}, \sqrt{1-p/2})$, $\Gamma_1^{(B)} = I \otimes \mathrm{diag}(\sqrt{p/2}, -\sqrt{p/2})$, where $p = 1 - \exp(-\gamma t)$, γ is the phase damping rate [28, 29].

Let $\varepsilon(\cdot)$ represent the operator of decoherence. Then under the phase flip channel we have

$$\begin{aligned}\varepsilon(\rho) \;=\; & \frac{1}{4}(I\otimes I + I\otimes s\sigma_3 + (1-p)^2 c_1\sigma_1\otimes\sigma_1 \\ & +(1-p)^2 c_2\sigma_2\otimes\sigma_2 + c_3\sigma_3\otimes\sigma_3). \end{aligned} \tag{41}$$

As $\varepsilon(\rho)$ satisfy conditions in Eq.(27), (28), and the quantum deficit of the ρ^{ab} under the phase flip channel is given by

$$\begin{aligned}\Delta^{\rightarrow}(\varepsilon(\rho^{ab})) \;=\; & \frac{1}{4}[(1-c_3+\sqrt{s^2+(1-p)^4(c_1+c_2)^2})\log(1-c_3+\sqrt{s^2+(1-p)^4(c_1+c_2)^2}) \\ & +(1-c_3-\sqrt{s^2+(1-p)^4(c_1+c_2)^2})\log(1-c_3-\sqrt{s^2+(1-p)^4(c_1+c_2)^2}) \\ & +(1+c_3+\sqrt{s^2+(1-p)^4(c_1+c_2)^2})\log(1+c_3+\sqrt{s^2+(1-p)^4(c_1+c_2)^2}) \\ & +(1+c_3-\sqrt{s^2+(1-p)^4(c_1+c_2)^2})\log(1+c_3-\sqrt{s^2+(1-p)^4(c_1+c_2)^2})] \\ & -\frac{1}{4}[(1+s-c_3)\log(1+s-c_3)+(1+s+c_3)\log(1+s+c_3) \\ & +(1-s-c_3)\log(1-s-c_3)+(1-s+c_3)\log(1-s+c_3)]. \end{aligned} \tag{42}$$

As an example, for $s=0.3, c_1=0.3, c_2=-0.4, c_3=0.56$, the dynamic behavior of correlation of the state under the phase flip channel is depicted in Fig.2. We find that the concurrence C is greater than the deficit $\Delta^{\rightarrow}$ for $0\le P\le 0.237211$ and a sudden death of entanglement appears at $p=0.321904$, here one sees that the concurrence become zero after the transition. Therefore for these states the entanglement is weaker against the decoherence than the quantum deficit.



FIG. 2: Concurrence(blue dashed line) and quantum deficit(red solid line) under phase flip channel for $s=0.3$, $c_1=0.3$, $c_2=-0.4$ and $c_3=0.56$.

V. SUMMARY

We have studied the correlation for a class of X states. The level surfaces of quantum deficit have been depicted. For $r=s=0$ our results reduce to the ones for Bell-diagonal states. For nonzero s, it has been shown that the level surfaces of quantum deficit may have quite different geometry and topology. The quantum deficit become smaller in certain time interval for some initial states, and for some states the entanglement is weaker against the decoherence than the deficit.

Acknowledgments This work was supported by the NSFC 11101017, KZ201210028032 and

PHR201007107.

---

[1] C. H. Bennett, D. P. DiVincenzo, C. A. Fuchs, T. Mor, E. Rains, P. W. Shor, J. A. Smolin and W. K. Wootters, Phys. Rev. A **59**, 1070 (1999).
[2] W. H. Zurek, Ann. Phys.(Leipzip)9, 5(2000); L. Henderson and V. Vedral, J. Phys. A **34**, 6899 (2001); H. Ollivier and W. H. Zurek, Phys. Rev. Lett. **88**, 017901(2002).
[3] B. Li, Z. X.Wang and S. M. Fei, Phys. Rev. A **83**, 022321 (2011).
[4] Q. Chen, C. Zhang, S. Yu, X. X. Yi and C. H. Oh, Phys. Rev. A **84**, 042313 (2011).
[5] M. Shi, C. Sun, F. Jiang, X. Yan and J. Du, Phys. Rev. A **85**, 064104 (2012).
[6] M. D. Lang and C. M. Caves, Phys. Rev. Lett. **105**, 150501 (2010).
[7] S. Luo and S. Fu, Phys. Rev. A **82**, 034302 (2010).
[8] B. Dakic, V. Vedral and C. Brukner, Phys. Rev. Lett. **105**, 190502 (2010).
[9] G. Passante, O. Moussa and R. Laflamme Phys. Rev. A **85**, 032325 (2012).
[10] T. Werlang, S. Souza, F. F. Fanchini and C. J. Villas Boas, Phys. Rev. A **80**, 024103 (2009); B. Wang, Z. -Y. Xu, Z. -Q. Chen and M. Feng, Phys. Rev. A **81**, 014101 (2010); L. Mazzola, J. Piilo and S. Maniscalco, Int. J. Quan. Inf. **9**, 981 (2011); R. Auccaise, L. C. Céleri, D. O. Soares-Pinto, E. R. de Azevedo, J. Maziero, A. M. Souza, T. J. Bonagamba, R. S. Sarthour, I. S. Oliveira and R. M. Serra, Phys. Rev. Lett. **107**, 140403(2011).
[11] J. Oppenheim, M. Horodecki, P. Horodecki and R. Horodecki, Phys. Rev. Lett. **89**, 180402 (2002).
[12] M. Horodecki, K. Horodecki, P. Horodecki, R. Horodecki, J. Oppenheim, A. Sen De and U. Sen, Phys. Rev. Lett. **90**, 100402 (2003).
[13] S. Luo, Phys. Rev. A **77**, 042303 (2008).
[14] M. Piani, P. Horodecki and R. Horodecki, Phys. Rev. Lett. **100**, 090502 (2008).
[15] S. Wu, U. V. Poulsen and K. Mølmer, Phys. Rev. A **80**, 032319 (2009).
[16] K. Modi, T. Paterek, W. Son, V. Vedral and M. Williamson, Phys. Rev. Lett. **104**, 080501 (2010).
[17] G. Adesso and A. Datta, Phys. Rev. Lett. **105**, 030501 (2010).
[18] P. Giorda and M. G. A. Paris, Phys. Rev. Lett. **105**, 020503 (2010).
[19] K. Modi, A. Brodutch, H. Cable, T. Paterek and V. Vedral, arXiv: 1112.6238 (2011).
[20] W. H. Zurek, Phys. Rev. A **67**, 012320 (2003).
[21] M. Horodecki, P. Horodecki, R. Horodecki, J. Oppenheim, A. Sen De, U. Sen and B. Synak, Phys. Rev. A **71**, 062307 (2005); M. Horodecki, P. Horodecki and J. Oppenheim, Phys. Rev. A **67**, 062104 (2003).
[22] B. Synak, K. Horodecki and M. Horodecki, J. Math. Phys. **46**, 082107 (2005).
[23] A. Streltsov, H. Kampermann and D. Bruß, Phys. Rev. Lett. **108**, 250501 (2012).
[24] T. K. Chuan, J. Maillard, K. Modi, T. Paterek, M. Paternostro and M. Piani, Phys. Rev. Lett. **109**, 070501 (2012).
[25] A. Streltsov, H. Kampermann and D. Bruß, Phys. Rev. Lett. **106**, 160401 (2011).
[26] H. Kim, M. R. Hwang, E. Jung and D. Park, Phys. Rev. A **81**, 052325 (2010).
[27] M. A. Nielsen and I. L. Chuang, *Quantum Computation and Quantum Information* (Cambridge University Press, Cambridge, UK, 2000).
[28] J. Maziero, L. C. Céleri, R. M. Serra and V. Vedral, Phys. Rev. A **80**, 044102 (2009).
[29] T. Yu and J. H. Eberly, Phys. Rev. Lett. **97**, 140403 (2006).